

Diffuse x-rays: Directly Observing Dark Matter?

Michael McNeil Forbes

Nuclear Theory Group, Department of Physics, University of Washington, Seattle, WA 98195-1560, USA

Ariel R. Zhitnitsky

The University of British Columbia, Department of Physics and Astronomy, Vancouver, B.C., V6T 1Z1, Canada

Several independent observations of the Galactic core suggest hitherto unexplained sources of energy. We suggest that dark matter in the form of dense antimatter nuggets could provide a natural site for electron and proton annihilation, providing 511 keV photons, gamma-rays, and diffuse keV x-ray radiation. We show that identifying dark matter as antimatter nuggets is consistent with the observed emissions, and we make definite predictions about their spectrum and morphology. If correct, our proposal not only identifies dark matter and explains baryogenesis, but allows x-ray observations to directly probe the matter distribution in our Galaxy.

PACS numbers: 95.35.+d, 98.70.-f, 12.38.-t

I. INTRODUCTION:

Two of the outstanding cosmological mysteries—the natures of dark matter and baryogenesis—might be explained by the idea that dark matter consists of Compact Composite Objects (CCOs) [1, 2, 3], similar to Witten’s strangelets [4]. The idea is that these CCOs—nuggets of dense matter and antimatter—form at the same QCD phase transition as conventional baryons (neutrons and protons), providing a natural explanation for the similar scales $\Omega_{\text{DM}} \approx 5\Omega_{\text{B}}$. Baryogenesis proceeds through a charge separation mechanism: both matter and antimatter nuggets form, but the natural CP violation of the so-called θ term in QCD¹ drives the formation of more antimatter nuggets than matter nuggets, resulting in the left-over baryonic matter that forms visible matter today (see [2] for details). The idea that CP violation may be able to effectively drive charge separation may already have found experimental support through the Relativistic Heavy Ion Collider (RHIC) at Brookhaven [5].

This mechanism requires no fundamental baryon asymmetry to explain the observed matter/antimatter asymmetry: $B_{\text{univ}} = B_{\text{DM}} + B_{\text{Visible}} - \bar{B}_{\text{DM}} = 0$ where $B_{\text{DM}}(\bar{B}_{\text{DM}})$ is the total (anti)baryon charge contained in the dark (anti)matter nuggets. From this and the observed relation $\Omega_{\text{DM}} \approx 5\Omega_{\text{B}}$ we have the approximate ratios $\bar{B}_{\text{DM}}:B_{\text{DM}}:B_{\text{Visible}} \simeq 3:2:1$.

Unlike conventional dark matter candidates, dark antimatter nuggets would be strongly interacting, but macroscopically large objects, and would not contradict any of

the many known observational constraints on dark matter or antimatter [3, 6] for three reasons:

- (1) They carry a huge (anti)baryon charge $|B| \approx 10^{20}\text{--}10^{33}$, so they have an extremely tiny number density. This would explain why they have not been directly observed on earth. The local number density of dark matter particles with these masses is small enough that interactions with detectors are exceedingly rare and fall within all known detector and seismic constraints [3].
- (2) They have nuclear densities, so their interaction cross-section is small $\sigma/M \approx 10^{-13}\text{--}10^{-9} \text{ cm}^2/\text{g}$. This is well below typical astrophysical and cosmological limits which are on the order of $\sigma/M < 1 \text{ cm}^2/\text{g}$. Dark-matter–dark-matter interactions between these nuggets are thus negligible.
- (3) They have a large binding energy such that baryons in the nuggets are not available to participate in big bang nucleosynthesis (BBN) at $T \approx 1 \text{ MeV}$. In particular, we suspect that the baryons in these nuggets form a superfluid with a gap of the order $\Delta \approx 100 \text{ MeV}$. This scale would provide a natural explanation for the observed photon to baryon $n_B/n_\gamma \sim 10^{-10}$, which requires that formation of the nuggets stop at precisely $T_{\text{formation}} = 41 \text{ MeV}$ [2, 7]. At temperatures below this, incident baryons with energies below the gap would Andreev reflect rather than become incorporated into the nugget.²

Thus, on large scales, the nuggets would be sufficiently dilute that they would behave as standard collisionless cold dark matter (CDM). When the number densities of both dark and visible matter become sufficiently high, however, dark-antimatter–visible-matter collisions may release significant radiation and energy.

¹ If θ is non-zero, one must confront the so-called strong CP problem whereby some mechanism must be found to make the effective θ parameter extremely small today in accordance with measurements. One of the most natural resolutions is through a dynamical axion. Domain walls associated with this field (or ultimately, whatever mechanism resolves the strong CP problem) play an important role in forming these nuggets, and may play an important role in their ultimate stability. See [1, 2] for details.

² If this mechanism is verified, then this value of $T_{\text{formation}}$ would provide one of the most accurate constraints on the properties of high-density QCD matter.

In this paper, we explain how the interaction between visible matter and dark antimatter nuggets in our Galaxy could naturally resolve several outstanding observational puzzles. These annihilations in the galactic core may be the primary source of detected 511 keV radiation, and the primary source of energy sustaining diffuse keV emissions. We consider three independent observations of diffuse radiations from the galactic core:

- (1) SPI/INTEGRAL observes 511 keV photons from positronium decay that is difficult to explain with conventional astrophysical positron sources [8, 9, 10]. Dark antimatter nuggets would provide an unlimited source of positrons as suggested in [11, 12].
- (2) *Chandra* observes a diffuse keV x-ray emission that greatly exceeds the energy from identified sources [13]. Visible-matter/dark-antimatter annihilation would provide this energy.
- (3) EGRET/CGRO detects MeV to GeV gamma-rays, constraining antimatter annihilation rates.

The main goal of this paper is to demonstrate that these observations suggest a common origin for both the 511 keV radiation and the diffuse keV x-ray emission. Assuming dark antimatter nuggets to be the common source, we extract some phenomenological parameters describing their properties. We show that dark antimatter is consistent with all of these data, and that it may fully explain the missing sources of emission without requiring the invention of new non-baryonic fields for dark matter. Finally, we argue that the spectrum of the keV emission should be independent of position, and that there should be a spatial correlation between the various emissions and the distributions of dark and visible matter. These concrete predictions should allow our proposal to be verified or ruled out by observations and analysis in the near future.

Our purpose here is not to discuss details of a particular model for the nuggets, but to show generally that macroscopic antimatter nuggets could easily explain the missing energy and missing positrons in the core of our Galaxy. To this end, we postulate a few basic properties to be discussed in detail elsewhere [6]:

- (A.1) The antimatter nuggets provide a virtually unlimited source of positrons (e^+) such that impinging electrons (e^-) will readily annihilate at their surface through the formation of positronium [11, 12].

About a quarter of the positronium annihilations release back-to-back 511 keV photons. If this reaction occurs on the surface, on average one of these photons will be absorbed by the nugget while the other will be released.

- (A.2) The nuggets provide a significant source of anti-baryonic matter such that impinging protons will annihilate. We assume that the proton annihilation rate is directly related to that of electrons through a suppression factor $f < 1$ to be discussed below.

Proton annihilation events will release about $2m_p \approx 2$ GeV of energy per event and will occur close to the surface of the nugget creating a hot spot that will mainly

radiate x-ray photons with keV energies rather than GeV gamma-rays, (see the appendix for a detailed discussion.) To connect these emissions with observations, we make an additional assumption:

- (A.3) We assume that these emitted 511 keV photons dominate the observed 511 keV flux and that these proton annihilations provide the dominant source of energy fueling the keV emission.

This allows us to use the 511 keV observations to estimate the electron annihilation rate and the *Chandra* flux to estimate the proton annihilation rate. This assumption may not be fully realized—the observed fluxes may be a combination of various conventional and unconventional sources—but by assuming that the majority of the flux originates from matter-antimatter annihilations, we are able to provide constraints on the annihilation properties of the nuggets. As we shall discuss, these constraints, although only bounds, are non-trivial and could have ruled out our proposal.

II. PROPOSAL:

The primary feature of our proposal is that the antimatter nuggets provide a single annihilation target for both electrons and protons. As a result, both the 511 keV emission from electron annihilation and the x-ray emission from proton annihilation should originate from the same regions of space, with the local rates of annihilation proportional to the product of the number densities of visible and dark matter $n_V(\vec{r})n_D(\vec{r})$.

By comparing several different observational sources, one may remove the dependence on the dark and visible matter distributions because all observations integrate the radiation along the same line of sight from the earth to the core of the Galaxy. We make the approximation that the relevant line-of-sight average is the same for the three sets of data we consider. As a result, direct comparisons between the data provide non-trivial insights about the properties of dark antimatter, independent of the matter distributions.

Chandra: X-ray emissions from the galactic core provide a puzzling picture: they seem to indicate that an 8 keV thermal plasma is being maintained, but the source of energy fueling this plasma is a mystery. After subtracting known x-ray sources from the *Chandra* x-ray images of the galactic core, one finds a residual diffuse thermal x-ray emission with a thermal component well described by a hot 8 keV plasma with surface brightness $\Phi_{Chandra} = (1.8\text{--}3.1) \times 10^{-6}$ erg/cm²/s/sr [13]. To sustain such a plasma would require some 10^{40} erg/s of energy in the galactic core, which is much more than the observed rate of supernovae, for example, can explain [13].

Protons annihilating with dark antimatter would release some $2m_p \approx 2$ GeV of energy per event. Assuming

(A.3.) this to be the dominant source of energy sustaining the keV emission, one may relate the x-ray luminosity to the number of annihilations events observed along the line of sight. Proton/dark-antimatter annihilations in the galactic core must thus provide an energy input of

$$\Phi_{Chandra} \approx 2 \times 10^{-6} \frac{\text{erg}}{\text{cm}^2 \cdot \text{s} \cdot \text{sr}} \quad (1)$$

into keV photons along our line of sight as observed in [13]. We shall see that our proposal can easily accommodate this flux.

SPI/INTEGRAL: An observed flux of 511 keV photons from the core of the Galaxy is another puzzle. It is clear that this comes from low-energy $e^-e^+ \rightarrow 2\gamma$ annihilation events (positronium decay), but the source of positrons is a mystery (the interstellar medium contains many electrons). After accounting for known positron sources, only a small fraction of the emission may be explained [8, 9, 10]. SPI has measured the flux of 511 keV photons from the galactic bulge to be $9.9_{-2.1}^{+4.7} \times 10^{-4}$ ph/cm²/s with a half maximum at 9° [8, 14, 15, 16].

As discussed in [11, 12], dark antimatter nuggets provide a natural source of positrons in the “atmosphere” of positrons (electrosphere) surrounding the nuggets that will readily annihilate with electrons from the ISM, (see equation (A1) in the appendix). The resulting spectrum will be dominated by the formation and subsequent decay of positronium, one quarter of which will decay from the $^1S_0$ state giving rise to two back-to-back 511 keV photons. Under assumption A.1., on average, one of these photons will be radiated. These $^1S_0$ annihilations account for approximately one quarter of the annihilations events: the rest occur through the $^3S_1$ state and give rise to a continuum below 511 keV that has also been measured [17], confirming the source as positronium decay.

Assuming (A.3.) that this e^-e^+ annihilation is the dominant source of the measured 511 keV photons, the flux from the core of the Galaxy provides an estimate of the rate of annihilation events. Unfortunately, the angular resolution of INTEGRAL is not very precise. To estimate the core flux requires some assumption about the spatial distribution of the annihilations. We simply use the estimates in [18, 19] to obtain an order of magnitude for same region observed by *Chandra* (1). Multiplying by a factor of 4 to account for the continuum emission, the rate of annihilations integrated along our line of sight is

$$\phi_{e^-e^+} \approx 10^{-1} \frac{\text{event}}{\text{cm}^2 \cdot \text{s} \cdot \text{sr}}. \quad (2)$$

EGRET: Some fraction of the proton annihilation events will produce high energy photons that may directly escape. These photons, with energies up to 1 GeV, should be detected by EGRET aboard the CGRO satellite, which has measured the intensity of photons from 30 MeV to 4 GeV from a central core region of 60° longitude and 10° latitude [20, 21]. Much of this emission

is due to cosmic rays interacting with visible matter. We therefore define the parameter $a_{\text{EGRET}} < 1$ to be the fraction of the total EGRET intensity due to direct photon production from proton/dark-antimatter annihilations.

We may thus use the EGRET data to provide two constraints on proton/dark antimatter annihilations. Performing the energy integrals from 100 MeV to 1 GeV (the typical range of photon energy from p^+p^- annihilations) on the data from [21], we obtain the following integrated count $\phi_{p^+p^- \rightarrow \gamma}$ and energy Φ_{EGRET} surface brightnesses extracted from the EGRET data:

$$\phi_{p^+p^- \rightarrow \gamma} \approx 10^{-4} \frac{\text{ph}}{\text{cm}^2 \cdot \text{s} \cdot \text{sr}}, \quad (3a)$$

$$\Phi_{\text{EGRET}} \approx 10^{-7} \frac{\text{erg}}{\text{cm}^2 \cdot \text{s} \cdot \text{sr}}. \quad (3b)$$

III. COMPARISONS:

We now can compare (1), (2), and (3) to extract some properties of the visible matter/dark-antimatter annihilation rates.

A. *Chandra* Constraints:

Proton-antiproton annihilation should proceed at a rate $\phi_{p^+p^-} = f\phi_{e^-e^+}$ proportional to, but different from, that for electron-positron annihilation. The suppression factor f is due to several effects.

- (1) The most important is that the transition from quark-matter to vacuum occurs on a QCD scale and is very sharp: $\sim \text{fm} \sim (200 \text{ MeV})^{-1}$. This needs to be compared with the long de-Broglie wavelength of the incoming protons, which have momenta on the scale $p = m_p v \sim \text{MeV}$. As such, the effective potential for the incoming protons due to this transition will behave qualitatively as a delta-function and the reflection probability will be high, irrespective of whether the interaction is attractive or repulsive. (This is a standard and general result from one-dimensional quantum mechanics.)
- (2) Additional suppression may result from superfluid correlations in the nugget. This would lead to the analogue of Andreev reflection in superconductors where incident protons reflect, rather than penetrate.
- (3) Finally, charge exchange reactions may take places which allow the nuggets to maintain overall neutrality charge neutrality, despite the higher rate of positron annihilation than proton annihilation.

Together these effects lead to a suppression factor f which, in principle, may be calculated. The result, however, will be highly model dependent. Instead, we introduce this as a phenomenological parameter. Note that this parameter must be less than unity: $f < 1$. If observations determine this parameter to be large, the model as a whole can be ruled out. Thus, even without a precise

calculation of f , observations provide a nontrivial test of our proposal. Once f is known, one can estimate the rate at which energy is deposited into the core of the Galaxy.

Our picture is that the fraction f of protons that annihilate penetrate a short way into the dark antimatter nuggets before annihilation. Annihilation produces a handful of high energy particles, (mostly Goldstone bosons). About half of these penetrate deep within the nuggets, ultimately thermalizing and radiating low-energy photons (this emission is not subject of the present work) but the remaining half stream toward the surface of the nugget, ultimately cascading into hundreds of highly energetic positrons that stream into the electrosphere. These positrons are accelerated by the strong electromagnetic fields present at the surface of the nuggets and radiate some fraction g of the total 2 GeV annihilation energy through Bremsstrahlung radiation with a typical keV energy scale set by the plasma frequency of the photons in the electrosphere.³

A detailed account of this process is presented in the appendix. Note: we expect that this mechanism of radiation—where energetic electrons are accelerated by a strong electric field present in the electrosphere—is also the dominant mechanism for radiation in strange quark stars, yet it appears to have been overlooked in the literature of radiation from the surface of strange quark stars [22].

The end result is that each proton annihilation produces a “hot spot” on the surface of the nugget that emits a broad spectrum of photons on the 10 keV scale containing a total energy of $g \times 2$ GeV. Although the details of this spectrum are difficult to calculate and are model dependent, (see the discussion in the appendix), one immediately has a strong prediction that the spectrum should be relatively independent of the location of the emission. This is in marked contrast with thermal plasma emission where the spectrum depends sharply on the temperature of the plasma, which in turn depends strongly on local parameters, such as the rate of energy input and particle density in the surrounding environment.⁴ We note that this is already consistent with analyses of the excess EGRET emission [23].

Our proposal would thus resolve the problem of heating an 8 keV thermal plasma in the core of the Galaxy

by replacing the plasma with an emitting hot-spot on the antimatter nugget surface fueled by proton annihilation. The similarity to a thermal spectrum arises from Bremsstrahlung emission in the electrosphere of the nuggets stimulated by the strong electric fields present near the surface of quark matter objects systems [24, 25, 26, 27]. This is discussed further in section A 5 of the appendix.

The x-rays detected by *Chandra* allow us to directly compare the e^-e^+ annihilation rate (2) with the observed keV emission $\Phi_{Chandra} = (2 \text{ GeV}) f g \phi_{e^-e^+}$. From this we may estimate the total suppression fraction $f g$ from (1) and (2):

$$f g = \frac{\Phi_{Chandra}}{(2 \text{ GeV}) \phi_{e^-e^+}} \approx 6 \times 10^{-3}. \quad (4)$$

As discussed in the appendix, we expect the suppression factor g to be on the order of $\frac{1}{10} < g < \frac{1}{2}$. The remaining suppression factor f could easily be accounted for by strong interface effects at the surface of dark antimatter nuggets as mentioned above. The parameter f , however, is not entirely free: It would be difficult, for example, to explain a value of $f > 1$ greater than unity due to the suppression factors discussed above. Thus, (4) provides a nontrivial test of our proposal that could rule it out if strong arguments could be given that f must be greater than unity. In principle, f and g are calculable from detailed models of the nuggets, providing a direct test of any quantitative model for dark antimatter.

B. EGRET Constraints

We now consider the EGRET constraints. If proton annihilations occur inside of the antimatter nuggets, then most of the decay products will be strongly interacting and will scatter into positrons or thermalize as discussed above. Occasionally, however, the annihilation will directly produce a high-energy photon with an energy of 100 MeV to 1 GeV that can easily escape the nugget. These processes must be compared with similar processes involving gluons rather than photons. Direct photon production is thus suppressed by a factor of $\alpha/\alpha_S \sim 1/40$ – $1/100$ due to the relative weakness of the electromagnetic interaction compared to the strong interaction. Thus, we may convert the rate of proton annihilations to a rate of direct photon emission along our line of sight, and compare the integrated EGRET data to the proton-antiproton annihilation rates $a_{EGRET} \phi_{p^+p^- \rightarrow \gamma} = (\alpha/\alpha_S) \Phi_{Chandra}/(2 \text{ GeV}) = f (\alpha/\alpha_S) \phi_{e^-e^+}$. Using (1) and (3a) we obtain

$$a_{EGRET} = \frac{\alpha}{\alpha_S} \frac{\Phi_{Chandra}/(2 \text{ GeV})}{\phi_{p^+p^- \rightarrow \gamma}} \approx 10^{-1}, \quad (5)$$

where a_{EGRET} is the fraction of the total EGRET intensity resulting from this direct photon emission. (Note: this estimate does not include photons from neutral meson

³ The factor $1 - g$ is introduced to account for the fraction of the 2 GeV annihilation energy not detected by *Chandra*. This includes the half of the energy lost geometrically deep into the nugget producing very low energy photons, as well as the radiated energy in the form of very hard x-rays (> 10 keV) not detected by *Chandra*. Such hard x-rays may in principle be emitted by this mechanism as discussed in the appendix. It also accounts for other energy losses, such as neutrino emissions, but these do not play a dominant role.

⁴ There will be a mild dependence of the spectrum on the thermal distribution of the incident protons, but this is a small effect as typical interstellar temperatures are much below the 10 keV scale of these emissions.

decays such as $\pi \rightarrow 2\gamma$ because these photons cannot escape the nuggets. This is discussed in detail in A3.) Finally, we may compare the amount of energy released as high energy photons with the integrated EGRET energy flux $a_{\text{EGRET}} \Phi_{\text{EGRET}} = (\alpha/\alpha_S) \Phi_{\text{Chandra}}$ to obtain an independent estimate of a_{EGRET} from (1) and (3b)

$$a_{\text{EGRET}} = \frac{\alpha}{\alpha_S} \frac{\Phi_{\text{Chandra}}}{\Phi_{\text{EGRET}}} \approx 3 \times 10^{-1}. \quad (6)$$

This is consistent with (5) to within the errors of our approximations, and demonstrates the importance of the GeV energy scale for both the x-rays band measure by *Chandra* and the gamma ray band measured by EGRET. This “accidental” coincidence is naturally provided by the QCD scale in our proposal. Together, equations (5) and (6) suggest that visible-matter/dark-antimatter annihilations could account for a small fraction ($\sim 10\%$) of the measured EGRET intensity. Much of the EGRET intensity has been accounted for, and future improvements in this accounting could provide a tight constraint on our proposal. It is nontrivial that the measured fraction $a_{\text{EGRET}} < 1$ be less than one: If the observations showed a_{EGRET} to be greater than unity, then one would have had a strong case against our proposal.⁵

We have made several assumptions in our estimates that must be cleaned up in order to make quantitative predictions. We have assumed that the line-of-sight integral over the matter distribution is the same for each observation, and that the phenomenological suppression factors f , g , as well as the fraction a_{EGRET} , are independent of the local environment and hence factor from this integral. In reality these factors will depend on the energies of the events, the nature of the environment (ionized vs. neutral), the local velocities of the visible and dark matter, and a host of similar factors. We may lump these factor together for any given line-of-sight, but it will be important to quantify how these factors vary as different line-of-sight averages are performed in order to make quantitative comparisons along different lines of sight.

We have also purposefully neglected any particular models of the matter distribution in our Galaxy. By comparing several observations along similar lines of sight, the results presented in this work do not depend on details such as the typical baryon charge B of the nuggets, nor do they depend on the exact matter distributions. Postulating a distribution leads to a direct connection

between the observed fluxes and properties of the dark antimatter nuggets such as the interaction cross-section $\sigma/M \approx 10^{-13}\text{--}10^{-9} \text{ cm}^2/\text{g}$. This is consistent with all known observations and interaction constraints, but a more careful analysis could significantly narrow the window of parameters. We shall discuss these and other details about the nature of dark antimatter elsewhere [6].

IV. PREDICTIONS:

We have shown that the observed excess 511 keV photons and thermal x-ray radiation from the core of our Galaxy may be directly explained by the hypothesis that much of the dark matter in the core of our Galaxy is in the form of dark antimatter nuggets. In this case, both observations have the same physical origin and are naturally correlated. The observations suggest very reasonable values for f and g —the suppression of the proton to electron annihilation rates and fraction of energy emitted—and a_{EGRET} —the fraction of EGRET photons arising from dark matter annihilations. These parameters satisfy the nontrivial constraints $f < 1$ and $a_{\text{EGRET}} < 1$, either of which could rule out our proposal.

Our proposal also makes the definite prediction that the 511 keV flux and the thermal x-ray flux should be correlated. For example, *Chandra* has detected a diffuse x-ray emission with flux $6.5 \times 10^{-11} \text{ erg/cm}^2/\text{s/deg}^2$ from a region of the disk 28° off the center [28]. This is one order of magnitude smaller than (1), and so the integrated positronium annihilation flux should be similarly reduced from (2): $\phi_{e^-e^+} = 10^{-2} \text{ event/cm}^2/\text{s/sr}$. We predict that the local 511 keV flux should be one quarter of this rate.

These fluxes should also be correlated with the product of the distributions of dark and visible matter $n_D(\vec{r})n_V(\vec{r})$. The angular resolution of INTEGRAL is not spectacular, but sufficient that the different source distributions can be compared. For example, as discussed in [29], the measured angular distribution is consistent spatial distribution models of $n_V(\vec{r})n_D(\vec{r})$. Unlike the mechanism suggested in [29], however, our proposal can also easily account for the observed luminosity.

Checking the angular 511 keV distribution with the matter distribution inferred from the *Chandra* x-ray data should provide a nontrivial confirmation or refutation of this model. One also has the possibility of looking for correlations elsewhere from clusters or in places where dark matter halo clumps pass through the disk. The problem here becomes one of sensitivity: INTEGRAL is not sensitive enough to detect the 511 keV emissions from extragalactic sources. *Chandra*, however, may still be used to detect extra-galactic x-ray emission. If our proposal is correct, diffuse x-ray measurements with subtracted sources could be directly translated into measurements of the integrated $n_V n_D$ distribution. This could be further checked by comparing the inferred matter distributions with those measured by weak lensing, rotation curves, and other emissions.

⁵ There is still some debate about the fraction of the observed EGRET emission that can be “explained” with conventional backgrounds. For example, [23] suggests that the observations support an excess that may result from WIMP dark-matter annihilations, while [20] suggests that much of this could be explained by cosmic rays. Our point is that the present constraints still leave a large region to accommodate our proposal. In principle, however, as the backgrounds are better understood, they will place stronger and stronger constraints on a_{EGRET} , possibly ruling out our proposal.

Finally, this proposal makes definite testable predictions for the properties of the emitted spectra. In particular, both the 511 keV spectrum and the diffuse keV spectrum are produced on an event-by-event basis, and are thus independent of the rate at which the annihilation processes occur (the timescale for the emitting “hot spots” to radiate is much shorter than the local collisional frequency, so the nuggets do not “integrate” the effects of the annihilations.) The observed spectra should thus be largely *independent of the direction of observation*.⁶ Only the intensity should vary as a function of the collision rate, and this should be correlated with the visible/dark matter distribution as discussed above.

V. CONCLUSION:

Our dark matter proposal not only explains many astrophysical and cosmological puzzles, but makes definite predictions about the correlations of the dark and visible matter distributions $n_V n_D$ with 511 keV, x-ray and gamma-ray emissions. In addition, it makes the definite prediction that the spectra of the emissions should be vir-

tually independent of the local environment. Such correlations and spectral properties would be very difficult to account for with other dark matter candidates. Future observations may thus confirm or rule out this theory. If confirmed, it would provide a key for many cosmological and astrophysical secrets, and finally unlock nature of dark matter.

As an aside, we mention that the mechanism for producing the x-ray radiation discussed in the appendix is likely the dominant mechanism for radiation from the surface of bare strange quark stars, should they exist, though it seems to have been neglected in the literature.

Acknowledgments

The authors would like to thank Jeremy Heyl and Tom Quinn for useful discussions, John Beacom, Francesc Ferrer, and Andrew Strong for comments on the manuscript, the PITP program where this project was started, the US Department of Energy for support under Grant DE-FG02-97ER41014, and the Natural Sciences and Engineering Research Council of Canada.

-
- [1] A. R. Zhitnitsky, JCAP **10**, 010 (2003), [arXiv:hep-ph/0202161](#).
 - [2] D. H. Oaknin and A. Zhitnitsky, Phys. Rev. D **71**, 023519 (2005), [arXiv:hep-ph/0309086](#).
 - [3] A. Zhitnitsky, Phys. Rev. D **74**, 043515 (2006), [arXiv:astro-ph/0603064](#).
 - [4] E. Witten, Phys. Rev. D **30**, 272 (1984).
 - [5] D. Kharzeev and A. Zhitnitsky, Nucl. Phys. A **797**, 67 (2007), [arXiv:0706.1026](#) [hep-ph].
 - [6] M. M. Forbes and A. R. Zhitnitsky, in prep.
 - [7] E. W. Kolb and M. S. Turner, *The Early Universe* (Westview Press, 5500 Central Avenue, Boulder, Colorado, 80301, 1994).
 - [8] J. Knödseder *et al.*, Astron. Astrophys. **411**, L457 (2003), [arXiv:astro-ph/0309442](#).
 - [9] J. F. Beacom and H. Yüksel, Phys. Rev. Lett. **97**, 071102 (2006), [arXiv:astro-ph/0512411](#).
 - [10] H. Yüksel, Nucl. Phys. B Proc. Suppl. **173**, 83 (2006), [arXiv:astro-ph/0609139](#).
 - [11] D. H. Oaknin and A. R. Zhitnitsky, Phys. Rev. Lett. **94**, 101301 (2005), [arXiv:hep-ph/0406146](#).
 - [12] A. Zhitnitsky, Phys. Rev. D **76**, 103518 (2007), [arXiv:astro-ph/0607361](#).
 - [13] M. P. Muno *et al.*, Astrophys. J. **613**, 326 (2004), [arXiv:astro-ph/0402087](#).
 - [14] P. Jean *et al.*, Astron. Astrophys. **407**, L55 (2003), [arXiv:astro-ph/0309484](#).
 - [15] J. Knödseder *et al.*, Astron. Astrophys. **441**, 513 (2005), [arXiv:astro-ph/0506026](#).
 - [16] P. Jean *et al.*, Astron. Astrophys. **445**, 579 (2006), [arXiv:astro-ph/0509298](#).
 - [17] G. Weidenspointner *et al.*, Astron. Astrophys. **450**, 1013 (2006), [arXiv:astro-ph/0601673](#).
 - [18] C. Boehm, D. Hooper, J. Silk, M. Casse, and J. Paul, Phys. Rev. Lett. **92**, 101301 (2004), [arXiv:astro-ph/0309686](#).
 - [19] D. Hooper *et al.*, Phys. Rev. Lett. **93**, 161302 (2004), [arXiv:astro-ph/0311150](#).
 - [20] A. W. Strong, I. V. Moskalenko, and O. Reimer, Astrophys. J. **613**, 962 (2004), [arXiv:astro-ph/0406254](#).
 - [21] T. Kamae, T. Abe, and T. Koi, Astrophys. J. **620**, 244 (2005), [arXiv:astro-ph/0410617](#).
 - [22] D. Page, M. Prakash, and V. V. Usov, private communication, 2007.
 - [23] W. de Boer *et al.*, (2004), [arXiv:astro-ph/0408272](#).
 - [24] C. Alcock, E. Farhi, and A. Olinto, Astrophys. J. **310**, 261 (1986).
 - [25] C. Kettner, F. Weber, M. K. Weigel, and N. K. Glendenning, Phys. Rev. D **51**, 1440 (1995).
 - [26] J. Madsen, Phys. Rev. Lett. **87**, 172003 (2001), [arXiv:hep-ph/0108036](#).
 - [27] V. V. Usov, Phys. Rev. D **70**, 067301 (2004), [arXiv:astro-ph/0408217](#).
 - [28] K. Ebisawa *et al.*, Astrophys. J. **635**, 214 (2005), [arXiv:astro-ph/0507185](#).
 - [29] J. M. Freire *et al.*, (2006), [arXiv:hep-ph/0610240](#).
 - [30] D. T. Son and M. A. Stephanov, Phys. Rev. D **62**, 059902(E) (2000), [arXiv:hep-ph/0004095](#), Erratum to [31].
 - [31] D. T. Son and M. A. Stephanov, Phys. Rev. D **61**, 074012 (2000), [arXiv:hep-ph/9910491](#), Erratum: [30].
 - [32] K. Lawson and A. R. Zhitnitsky, JCAP **0801**, 022 (2008), [arXiv:0704.3064](#) [astro-ph].
 - [33] A. W. Steiner, S. Reddy, and M. Prakash, Phys. Rev. D **66**, 094007 (2002), [arXiv:hep-ph/0205201](#).
 - [34] V. V. Usov, T. Harko, and K. S. Cheng, Astrophys. J. **620**, 915 (2005), [arXiv:astro-ph/0410682](#).
 - [35] M. Alford and K. Rajagopal, (2006), [hep-ph/0606157](#).

- [36] E. Braaten, *ApJ* **392**, 70 (1992).
- [37] I. N. Mishustin, L. M. Satarov, T. J. Burvenich, H. Stoecker, and W. Greiner, *Phys. Rev.* **C71**, 035201 (2005), [arXiv:nucl-th/0404026](#).
- [38] J. D. Jackson, *Classical Electrodynamics*, Third ed. (John Wiley & Sons, Inc., New York, NY, 1999).

APPENDIX A: EMISSION OF KEV RADIATION

The goal of this appendix is to explain the process through which proton annihilation inside the dark anti-matter nuggets could release the keV scale photons observed *Chandra* emissions. The emission and spectrum of the 511 keV emission from electron-positron annihilation is discussed in detail in [11, 12, 32].

Although the details are somewhat complicated and model dependent, we shall argue that the characteristics of this emission are most strongly governed by the properties of the positron plasma with the emission scale being set by the plasma frequency near the surface of the nuggets. As such, the emission is largely insensitive to the model dependent details of the nugget core.

The basic idea is that protons could penetrate into the antimatter nuggets and annihilate near the surface. The 2 GeV of energy released by this annihilation is ultimately transferred to high-energy positrons that propagate out of the surface of the nuggets. These positrons are subsequently accelerated by the large electromagnetic fields present in the electrosphere, and release their energy as Bremsstrahlung radiation. The scale and spectral properties of this emission are determined by the properties of the electrosphere of the nuggets where the keV scale and approximately “thermal” nature of the spectrum arises quite naturally.

1. Structure of Nugget Surface

Here we briefly discuss the properties of the antimatter nuggets surface to set the scales. The radius of the nuggets depends on the mass, but must be larger than $R > 10^{-7}$ cm at the lower limit $|B| > 10^{20}$ set by terrestrial non-detection, but are most likely several orders of magnitude larger. The quark-matter core of the nuggets ends sharply on a fm scale as set by the nuclear scale. Near the surface, as the density falls, the quark matter will be charged due to the relatively large mass of the strange quark $m_s \sim 100$ MeV [24, 25, 26, 27]. (The matter may be charged throughout depending on the exact nature of the phase, but this does not affect our analysis.) Charge neutrality will be maintained through beta-equilibrium, which will establish a positron chemical potential $\mu_{e^+} \simeq 10$ MeV. (The precise value depends on specific details of quark matter phase and may range from a few MeV to hundreds of MeV, and is about an order of magnitude less than the quark chemical potential $\mu_q \simeq 10$ MeV [24, 33].) This will induce a thin but

macroscopic “electrosphere” of positrons surrounding the quark matter core in the transition region as $\mu_{e^+} \rightarrow 0$ in the vacuum.

The structure of this electrosphere has been considered for quark matter [24, 25] and the existence of this “transition region” is a very generic feature of these systems. It is the direct consequence of the Maxwell’s equations and chemical equilibrium. The region is called the electrosphere, emphasizing the fact that quarks and other strongly interacting particles are not present. In the case of antimatter nuggets the “electrosphere” comprises positrons.

The variation of this chemical potential $\mu_{e^+}(z)$, and the associated electric field $E(z)$ as a function of distance from the surface of the nugget z , may be computed using a mean-field treatment of the Maxwell equations [24, 25, 34]. For example, in the relativistic regime, one has [27]

$$\begin{aligned} \mu_{e^+}(z) &= \sqrt{\frac{3\pi}{2\alpha}} \frac{1}{(z+z_0)}, & z_0 &= \sqrt{\frac{3\pi}{2\alpha}} \frac{1}{\mu_0}, \\ E(z) &= \frac{1}{e} \frac{d\mu_{e^+}(z)}{dz}, \end{aligned} \quad (A1)$$

where $\mu_0 \equiv \mu_{e^+}(z=0) \sim 10$ MeV is the chemical potential realized in the nugget’s bulk. The corresponding results can be obtained outside of the relativistic regime, but they do not have a simple closed form.

Near the surface of the nugget, the typical scale is $z_0 \sim 10^{-11}$ cm ≈ 100 fm, thus, at a typical distance of $1000z_0 \sim 10^{-8}$ cm $\approx 10^5$ fm, the chemical potential will be of the keV scale. The electrosphere will extend well beyond this, but this is the most important scale for the present discussion.

2. Excitations and Interactions

The physics of radiation is dominated by the low-energy degrees of freedom present in the quark matter and in the electrosphere. Within the nuggets, the picture is of some charged phase of colour-superconducting matter with a gap of some 100 MeV, with a free Fermi-gas of positrons to maintain neutrality. (See [35] for a review of colour superconducting properties.)

Quarks and Gluons: The fundamental excitations in quark matter at sufficiently high densities are the quarks and gluons. These excitations are coloured and strongly interacting, but have “masses” on the order of the superconducting gap ~ 100 MeV. Thus, although the initial annihilation will produce these, the energy will ultimately be transferred to lower energy degrees of freedom.

Pseudo–Nambu–Goldstone Bosons (Mesons):

The spontaneous breaking of chiral symmetry by colour-superconductors gives rise to low-energy pseudo–Nambu–Goldstone modes with similar quantum numbers of mesons (pions, kaons etc.).

These objects, however, are collective excitations of the colour-superconducting state rather than vacuum excitations. The finite quark masses explicitly break the chiral symmetry, giving rise to these “pseudo”-Nambu-Goldstone modes on the order of 10 MeV (see [30, 31] for example). These modes are all strongly interacting and include both electrically charged ($K^\pm$, K , $\bar{K}$, $\pi^\pm$) and neutral (π^0 , η , η') modes.

Nambu-Goldstone Phonon: There is one truly massless Nambu-Goldstone mode associated with spontaneous baryon number violation. This mode, however, carries neither colour nor electric charge. Thus, it is extremely weakly coupled and plays an insignificant role in the radiative processes we consider here.

Positrons: The positrons exist as a Fermi liquid with chemical potential $\mu \sim 10$ MeV and Fermi energy p_F . ($p_F^2 = \mu^2 - m_{e^+}^2 \approx \mu^2$.) The total density of positrons is $n_{e^+}^{\text{tot}} = p_F^3/(3\pi^2)$, however, most of the states are Pauli blocked deep within the Fermi surface. For a given energy range $|E| < \omega$, we can estimate the number of available quasiparticle modes:⁷

$$n_{e^+} \simeq \frac{2\omega\mu p_F}{\pi^2} \approx \frac{2\omega\mu^2}{\pi^2}. \quad (\text{A2})$$

Photon: The electromagnetic interaction is mediated by a photon, but in the Fermi liquid, the interaction is screened and the photon has a plasma-frequency ω_p (see for example [24]):

$$\omega_p^2 \simeq \frac{4\alpha}{3\pi}\mu^2. \quad (\text{A3})$$

For $\mu_q \sim 100$ MeV, $\omega_p \approx 5$ MeV. This has several implications: 1) low energy photons do not propagate freely, 2) the $1/q^2$ singularity of the Coulomb interaction is softened $1/q^2 \rightarrow 1/(q^2 + \omega_p^2)$. Electromagnetic interactions are thus dominated by momenta transfer of $q \sim \omega_p$. We also consider the mean free path for high energy photons which can be estimated by considering the Compton scattering cross-section $\sigma_{e^+\gamma}$ in the positron’s rest frame (approximated for $\omega \gg m_{\text{eff}}$):

$$\sigma_{e^+\gamma} \simeq \frac{\pi\alpha^2}{\omega m_e} \ln\left(\frac{\omega}{m_e}\right), \quad l_\gamma^{-1} \simeq \sigma_{e^+\gamma} n_{e^+}(\omega_p). \quad (\text{A4})$$

From equations (A2) and (A3) we can estimate the mean-free path for charged particles in the positron gas. The cross-section between the charged meson modes (for example, the quark-matter equivalent of the π^+) and positrons is

$$\sigma_{e^+\pi^\pm} \simeq 2\pi \frac{\alpha^2}{\omega_p^2}, \quad l_{\pi^\pm}^{-1} \simeq \sigma_{e^+\pi^\pm} n_{e^+}(\omega_p). \quad (\text{A5})$$

The interaction here is dominated by momentum transfer of ω_p due to the screened photon. This has typical values of about $l_{\pi^\pm} \sim 10^{-9}$ cm for $\mu_{e^+} \sim 10$ MeV.

3. Energy Transfer

Now we consider how energy is transferred from proton annihilation events to radiation. Protons impinging upon dark antimatter nuggets will penetrate some characteristic depth $d \ll R$ much less than the radius of the nuggets. The depth will be set by the nuclear interaction scale: For example, if the annihilation p^+p^- cross-section is taken to be the vacuum value, the proton will have a lifetime of only 2 fm/c. It has been argued [37] that in nuclear matter, the lifetime could be considerably larger (possibly even 10–30 fm/c) due to the coherence required for the annihilation. The lifetime may be even longer in colour superconducting antimatter where the quantum numbers for all three anti-quarks must be correlated for a successful annihilation with incoming proton, perhaps even as large as 100 fm/c.

In any case, on the scale of the nuggets, the annihilations take place on the order of $h \sim 100$ fm or so from the surface, which is still within the regime where the quark matter is charged. Thus, the annihilations take place in a region of quark matter where there positrons are present to maintain beta-equilibrium and neutrality. These positrons are the lightest modes that couple strongly and will thus ultimately carry most of the annihilation energy as we shall now explain.

The proton annihilation will initially produce a handful of high energy gluons. These will stream away from the annihilation site. About half of these will stream towards the surface of the nuggets, and will ultimately transfer their energy to high-energy positrons. The other half, however, will proceed deep within the nuggets, and will most likely thermalize. Thus, only about half of the initial 2 GeV per annihilation will be converted into energetic positrons that propagate to the surface of the nuggets. The remaining energy will thermalize and subsequently be thermally radiated by the nuggets at very low temperatures. We account for this and other losses through a factor $g \lesssim 1/2$ which ultimately appears in our energy budget (4). We shall not further discuss this low temperature thermal radiation in this work, though it may have further observational consequences that need to be explored.

The remaining particles stream toward the surface of the nuggets and have a typical distance $d \sim \sqrt{2R\hbar} \sim$

⁷ In principle, we should include many-body corrections in these calculations. For example, the effective mass of the quasiparticles is $m_{\text{eff}} = m_e/2 + \sqrt{m_e^2/4 + \alpha\mu^2/(2\pi)}$ [36], but for $\mu = 10$ MeV, this is only a 30% correction. We neglect these corrections here for our order of magnitude estimates.

10^{-8} cm to travel before they hit the surface. At the annihilation point, with probability α/α_s , one of the decay products will be a high energy photon with $\omega \sim 1$ GeV $\gg \omega_p$. Such photons have a long mean-free paths, on the order of $l_\gamma^{1 \text{ GeV}} \sim 10^{-8}$ cm, and can directly leave the nuggets. These are the photons considered in (5) and (6) and should be observable by EGRET.

Only the photons directly produced by direct annihilation will have enough energy to escape. Photons from additional processes, such as the decay of the neutral pseudo-Nambu-Goldstone bosons ($\pi^0, \eta \rightarrow 2\gamma$), will have energies ~ 50 MeV. Not only are these processes unlikely to occur near the surface, (the typical width is $\Gamma \sim \alpha^2 m_{NG}^3 / (64\pi^3 f_\pi^2)$ corresponding to a huge mean free path⁸) but the mean free path for these photons is much shorter $l_\gamma^{50 \text{ MeV}} \sim 10^{-10}$ cm, so they will be absorbed long before they can escape. As a result, we do not include this energy in our estimates (5) and (6).

The remaining gluons will rapidly decay into the pseudo-Nambu-Goldstone collective modes, distributing the energy down to their mass scale of 10 MeV or so. As estimated in (A5), the charged modes scatter quite readily with positrons, transferring energies on the order of $\omega_p \sim 5$ MeV. These energetic positrons will travel a similar distance before scattering, (further distributing the energy down to this scale if it is above ω_p). Unlike the collective modes, however, the positrons are not bound to the nuggets.

Thus, through this mechanism, some fraction $g \lesssim 1/2$ of the initial 2 GeV annihilation energy will ultimately be transferred to positrons of energy $2 \sim 5$ MeV which stream out of the nuggets into the electrosphere. A small fraction α/α_s of the events may also produce GeV scale photons that directly escape. The rest of the energy is transferred deep into the nuggets, ultimately thermalizing and emitted at very low temperatures (to be discussed elsewhere [6]).

4. Bremsstrahlung Radiation

The remaining emission processes of the dark antimatter nuggets results from the physics of these 100 or so energetic positrons ($E \sim 5$ MeV) that enter the electrosphere of the nuggets. (For simplicity, we assume that all of the positrons have an energy of 5 MeV, being set by the plasma frequency of the photons that excited them. A detailed calculation of the spectrum would need to account for the scatter in this value.) As these energetic positrons move through the strong electric field $E(z)$,

they immediately start to emit photons. For a simple estimate of the timescale over which the energy is released, we treat these positrons classically. In this case, the total instantaneous radiated power is given by [38]

$$P_\gamma = \frac{2\alpha\gamma^6}{3} \left[(\dot{\vec{v}})^2 - (\vec{v} \times \dot{\vec{v}})^2 \right], \quad \gamma \equiv \frac{1}{\sqrt{1-v^2}}, \quad (\text{A6})$$

which we express in terms of the local electric $\vec{E}$ and magnetic $\vec{B}$ fields evaluated at the position of the particle [38],

$$P_\gamma = \frac{2\alpha^2\gamma^2}{3m^2} \left[(\vec{E} + \vec{v} \times \vec{B})^2 - (\vec{v} \cdot \vec{E})^2 \right]. \quad (\text{A7})$$

Now we can easily estimate the energy radiated per unit time by a positron in the background electric field (A1) as

$$P_\gamma \sim \mu^2 \cdot \frac{4\alpha^2\gamma^2}{9\pi} \cdot \left(\frac{\mu^2}{m^2} \right) \cdot \left(\frac{z_0}{z+z_0} \right)^4. \quad (\text{A8})$$

The total time for the positron with energy $\epsilon = 5$ MeV to release its energy in the form of synchrotron radiation is

$$\tau \sim \frac{\epsilon}{P_\gamma} \sim \frac{\epsilon}{\mu^2} \left(\frac{m^2}{\mu^2} \right) \frac{9\pi}{4\alpha^2\gamma^2} \left(\frac{z+z_0}{z_0} \right)^4. \quad (\text{A9})$$

Numerically, if we take $\mu \sim 10$ MeV and assume that most efficient radiation is happening at a typical z of a few times z_0 with an average $\gamma \sim \epsilon/(2m)$, we arrive at the estimate $c\tau \sim 10^{-8}$ cm.

In order for a photon to be efficiently emitted, its energy must be greater than the local plasma frequency ω_p . This is on the order of 5 MeV at the surface of the nugget, but rapidly falls off with $\mu_{e+}(z)$ so that by the time the photons have been formed at a distance on the order of 10^{-8} cm, the plasma frequency has dropped to the keV scale. This is consistent with fact that the wavelength of the emitted photons cannot be larger than length over which the photons are emitted $z = c\tau \sim 10^{-8}$ cm, corresponding to a minimum frequency $\omega \geq 2$ keV. This is not a lower bound on the spectrum, however, because additional scattering may also occur, distributing some of the initial energy of the positrons and extending the range of emission further into the electrosphere. This will soften the emission, but the characteristic energy scale will be keV.

5. Can Bremsstrahlung Mimic the Thermal Plasma Emission Spectra?

In this section, we show that the resulting keV Bremsstrahlung spectrum might be identified with the 8 keV thermal plasma emission spectrum used to fit the *Chandra* data. The most prominent feature of a thermal

⁸ Although the pseudo-Nambu-Goldstone modes have similar quantum numbers to the vacuum mesons, their masses differ by an order of magnitude from their vacuum counterparts. Coupled with the sharp quark-matter/vacuum interface, they thus cannot easily leave the quark matter, and will simply reflect off of the surface, ultimately re-scattering their energy into other particles.

plasma emission is the presence of the exponential factor which suppresses emission with $\omega \gg T$,

$$\frac{dP_\gamma(\omega)}{d\omega} \sim e^{-\omega/T}. \quad (\text{A10})$$

The characteristic spectrum is thus very broad, (almost flat up to a mild $\ln \omega$ dependence which has been omitted from (A10)) with a sharp cutoff at $\omega_c \sim T$. This exponential cut off is related to the minimum energy $mv_{\min}^2/2 \sim T$ required to emit a photon with frequency ω by thermal plasma. We note that the cutoff has not been directly observed because the detector band is limited at 8 keV, thus the spectrum might extend to higher energies (10 keV or so). (See [13] for details).

Although we cannot present a detail model-independent calculation of the spectrum, we shall argue that the qualitative features of the observed thermal spectrum are reproduced by field induced bremsstrahlung emission. In particular, there is a critical frequency ω_c above which emission is strongly suppressed, and the spectrum is very flat for low energies $\omega \ll \omega_c$. This behaviour is in sharp contrast with a black body spectrum, for example.

A quantitative treatment of the problem is difficult because the photon's emission must be calculated with a fully quantum mechanical treatment that takes into account the many-body effects and positron degeneracy.⁹ Ultimately this will be required to confirm or rule out any particular model. A qualitative picture for the spectrum, however, follows from simple arguments. Consider the motion of a relativistic classical particle in the background electric field (A1). The quantum mechanical emission of “soft” photons—photons whose energies are small compared with the typical interaction energy scale—is almost identical to that obtained by classical analysis (see for example [38]). Furthermore, the most important contribution comes from the small part of the classical trajectory with the largest instantaneous curvature. It is exactly this region that determines the critical value for ω_c and the qualitative features of the spectrum.

This key segment of the classical trajectory leads to the following emission spectrum [38],

$$\frac{dP_\gamma(\omega)}{d\omega} \sim \frac{\omega}{\omega_c} \int_{\omega/\omega_c}^{\infty} K_{5/3}(x) dx, \quad (\text{A11})$$

⁹ Photon emission in a dilute system will be suppressed by α^3 while equation (A7) explicitly demonstrates α^2 behavior. The enhancement is due to the coherent electric field (A1) which has $E \sim \frac{1}{e}$ behavior. This means that, in dense systems, the bremsstrahlung emission cannot be treated as a series of independent events (α^3 behavior), but must be treated as a coherent emission. The relevant parameter here is the ratio between the distance between electrons (μ^{-1}) and the typical energy transfer in each event (ω). The number of particles participating in the emission is very large, $((\mu/\omega)^3 \gg 1)$, therefore, the standard Bremsstrahlung emission technique cannot be used for our system. A precise calculation of the spectrum is beyond the scope of the present work.

with the asymptotic behaviors

$$\left. \frac{dP_\gamma(\omega)}{d\omega} \right|_{\omega \ll \omega_c} \sim \omega^{1/3}, \quad \left. \frac{dP_\gamma(\omega)}{d\omega} \right|_{\omega \gg \omega_c} \sim e^{-\omega/\omega_c}. \quad (\text{A12})$$

The main feature of the spectrum is that it is very broad and flat at frequencies below ω_c with a sharp cut off for $\omega \geq \omega_c$. In particular, when ω/ω_c changes by two orders of magnitude ($10^{-2} < \omega/\omega_c < 1$), the intensity changes only by factor 2. These features are very similar to those of a thermal plasma emission spectrum where the role of T is played by ω_c .

To estimate for ω_c , we note that the typical time scale for the changes in the system is $\delta t \sim \epsilon/eE$ where $\epsilon \sim \mu_0 + \epsilon_0$ is a total initial energy of the particle at $t = 0$. This parameter determines a curvature of the classical trajectory. The critical frequency of the emission ω_c in such a motion is $\omega_c \sim (\delta t)^{-1}$ can be estimated for the classical field (A1) as follows,

$$\omega_c \sim \frac{eE}{\mu_0 + \epsilon_0} \sim \sqrt{\frac{2\alpha}{3\pi}} \left(\frac{z_0}{z + z_0} \right)^2 \frac{\mu_0^2}{\mu_0 + \epsilon_0}. \quad (\text{A13})$$

Numerically this is on the order of 30 keV or so. Without a full quantum calculation, this is only an order of magnitude estimate, but it is consistent with our estimate for minimal frequency $\omega \geq 2$ keV obtained above.

We note here that this mechanism, whereby the moving positrons radiate as they are accelerated by the strong electric fields in the electrosphere, is likely the dominant mechanism radiating energy from the surface of strange quark matter such as strange stars. In this case, thermally excited electrons (strange stars would be matter rather than antimatter) would accelerate in the fields of the star's electrosphere, producing a similar radiation. This mechanism has apparently been overlooked in the literature [22]. Our estimates here provide a starting point, but a full quantum mechanical computation should be done to properly account for rate at which thermal energy may be radiated in these systems.

APPENDIX B: COSMIC RAYS IMPINGING ON NUGGETS

It has been suggested that much of the EGRET emission could result from cosmic rays (see for example [20]). This would affect the constraints on the a_{EGRET} as we have discussed, but raises another question: Could interactions between cosmic rays and nuggets have an observable consequence in the core of the Galaxy? The typical energy released by a cosmic-ray proton annihilating on a nugget would be 30 GeV rather than the 1 GeV released in the low-energy annihilations we have considered so far. In addition, there would be no suppression factor $f \sim 10^{-2}$ for these high energy events, thus, if there were a significant number of collisions, there would be an observable effect. As we shall see, the rate of collisions is very small,

so that the net energy released is four orders of magnitude less than our total budget, so we conclude that such processes may be neglected.

We start from analysis of the conventional and optimized models [20] for proton spectra as the most important portion of cosmic rays. Data presented in [20] suggests the cosmic-ray proton flux $I(E)$ to be about

$$E^2 \frac{dI(E)}{d\Omega dE} \sim 2 \times 10^{-1} \frac{\text{GeV}}{\text{cm}^2 \cdot \text{s} \cdot \text{sr}} \quad \text{where } 2 \text{ GeV} \leq E \leq 30 \text{ GeV}, \quad (\text{B1})$$

which corresponds to a total integrated flux

$$I = \int \frac{dI(E)}{d\Omega dE} d\Omega dE \sim 2 \times 10^{-1} \int \frac{dE}{E^2} \cdot \frac{\text{GeV} \cdot 4\pi}{\text{cm}^2 \cdot \text{s} \cdot \text{sr}} \sim 1 \frac{\text{proton}}{\text{cm}^2 \cdot \text{s}}. \quad (\text{B2})$$

The corresponding cosmic-ray proton flux (B2) can be used to estimate the total number of the collision events Φ_{CR} of cosmic-ray protons with the nuggets along the line of sight of the observation,

$$\Phi_{CR} \sim \int dr \int_{\Delta\Omega} d\Omega \cdot n_{DM}(l) \cdot I \cdot (4\pi R_0^2), \quad (\text{B3})$$

where $\Delta\Omega$ is the solid angle observed, and the integral $\int dr$ is performed over the line of sight of the observation. We assume that all cosmic-ray protons hitting the nuggets will annihilate, and that the total integrated flux of cosmic-ray protons, I —which enters equation (B3) and is estimated in (B2)—does not varies much with distance. (This would over-estimate the flux.)

The number density of the nuggets is approximately $n_{DM}(r) = \rho_{DM}(r)/M_{DM}$ with M_{DM} being a typical mass of the nugget. Numerically, Φ_{CR} is very small, about seven orders of magnitude smaller than the typical rate of collision between visible protons and nuggets. Such a suppression can be easily understood: When calculating the number of collision events between visible matter protons and nuggets along the same line of sight, one should replace I in formula (B3) by $n_B \cdot v$. Assuming $n_B \sim 1 \text{ cm}^{-3}$, and taking a typical $v/c \sim 10^{-3}$, one finds this seven orders of magnitude difference in the annihilation rate, which more than compensates for the enhanced annihilation rates (no factor of $f \sim 1/100$) and extra energy (30 GeV rather than 1 GeV) provided by these events.